arXiv:1304.5223v1 [math.RT] 18 Apr 2013

TWO-TERM TILTING COMPLEXES AND SIMPLE-MINDED SYSTEMS OF BRAUER STAR ALGEBRAS

AARON CHAN

Abstract. We show that any simple-minded system of Brauer star algebra is given by the restriction of the preimage of simple modules under a derived equivalence arising from two-term tilting complex concentrated in homological degree 0 and -1.

Keywords: simple-minded system, configuration, mutation, (two-term) tilting complex, Brauer tree.
MSC2010: 16G10, 18E30.

1. Introduction

Let A be a finite dimensional self-injective algebra over a field k. The structure of its stable module category $\underline{\text{mod}}$-A is in general difficult to study, mainly because it does not have objects which behave like projectives; instead one would use simple modules to study the homological properties of $\underline{\text{mod}}$-A. In [KL], Steffan Koenig and Yuming Liu proposed the notion for which a class of objects behaves like the set of simple objects, and called it *simple-minded system.* An analogue of such system which lives in the (bounded) derived category was introduced by Rickard [Ric1] and Al-Nofayee [AN]. This analogous idea is studied extensively by Koenig and Yang [KY] and is called *simple-minded collection* therein. In brief, Al-Nofayee showed that the class of simple-minded collections which are stable under the Nakayama functor is in one-to-one correspondence with the class of one-sided tilting complexes. It is well-known that the stable category $\underline{\text{mod}}$-A of finitely generated A-modules is a triangulated category, and this category is the same as the quotient $D^b(\text{mod-}A)/K^b(\text{proj-}A)$ of bounded derived category by the bounded homotopy category of complexes of projective A-modules, where the suspension functor of $\underline{\text{mod}}$-A is the inverse Heller translate Ω^{-1}. In particular, suppose $T \in K^b(\text{proj-}A)$ is a one-sided tilting complex and $E_T = \text{End}_{K^b(\text{proj-}A)}(T)$, then the derived equivalence $F_T = \text{Hom}(T, -) : D^b(\text{mod-}A) \to D^b(\text{mod-}E_T)$ restricts to a stable equivalence $\overline{F_T} : \underline{\text{mod}}\text{-}A \to \underline{\text{mod}}\text{-}E_T$. This is one of the main approaches to study the stable equivalences of self-injective algebras in the past two decades.

Moreover, using the results in [KY] and [KL], one can sees that restricting a Nakayama-stable simple-minded collection into the stable module category is in fact a simple-minded system, which revealed much more subtle connection between the derived categories and stable categories. In a recent collaboration [CKL] of the author with Koenig and Liu, we gave a detailed exposition on the simple-minded systems of representation-finite self-injective algebras, partly via such connection. One of the key results can be stated as follows: every simple-minded system of a representation-finite self-injective algebra arises as the image of the set of simple modules under some stable equivalence (of Morita type) induced by some derived equivalence. Moreover, this gives us a natural surjection from the class of tilting complexes to the class of simple-minded systems, which turns out to "respect mutation" as well. The present article aims to refine this connection in the case of Brauer star algebra.

By a result of Abe and Hoshino [AH], every derived equivalence given by tilting complex can be written as compositions of derived equivalence given by two-term tiling complexes (i.e. those that are concentrated in two consecutive homological degrees), and so we can think of two-term tilting complexes as "generators" for (one-sided) tilting complexes. On the other hand, in the emerging τ-tilting theory, the two-term tilting complexes of symmetric algebras are known to be in correspondence with support τ-tilting modules [AIR], which plays an important role in other areas of homological algebras, torsion pairs, 2-Calabi-Yau algebras, (cluster-)tilting and alike. To the best of our knowledge, the answer to the following problem concerning two-term tilting complex is still unknown to most self-injective algebras.

Problem 1.1. *Suppose A and B are derived equivalence algebras. Can we find a two-term tilting complex T concentrated in degree 0 and -1, so that $E_T = \mathrm{End}_{K^b(\text{proj-}A)}(T)$ is Morita equivalent to B?*

When A is a Brauer star algebra, the answer to this problem is well-known to experts in the field, which we will present in Section 3. To connect the study of two-term tilting complexes with that of simple-minded systems, we pose a strengthened version of the problem:

Problem 1.2. *Suppose A and B are self-injective k-algebras and $F : D^b(\text{mod-}A) \to D^b(\text{mod-}B)$ is a derived equivalence. Can we find a two-term tilting complex T concentrated in degree 0 and -1, so that the preimage of simple E_T-modules under the induced stable equivalence $\overline{F_T}$ coincide with the simple-minded system, which is given by the preimage of simple B-modules under the induced stable equivalence $\overline{F}$?*

This problem seems to be a difficult one in general, as all of the present article is devoted to solve this problem only in the case where A is a Brauer star algebra (symmetric Nakayama algebra). In the case when A is a representation-finite self-injective algebra, we know from the main results of [CKL] that positivity of this problem is equivalent to saying that the natural map from the class of two-term tilting complex to the class of simple-minded system is a surjection. For an algebra A, let 2-tilt(A) denote the set of two-term tilting complexes concentrated in degree 0 and -1 up to homotopy equivalence. The main result of the article is stated below; and it will be used to give a positive answer to Problem 1.2.

Theorem 1.3. *Let A be Brauer star algebra associated to the Brauer star with e edges and multiplicity $m > 1$. Suppose $T \in K^b(\text{proj-}A)$ is a two-term tilting complex, and $\overline{F_T} : \underline{\text{mod}}\text{-}A \to \underline{\text{mod}}\text{-}E_T$ be the induced stable equivalence. We have the following bijection:*

$$\begin{array}{rcl} \text{2-tilt}(A) & \leftrightarrow & \{\textit{simple-minded systems of } A\} \\ T & \mapsto & \overline{F_T}^{-1}(\textit{simple } E_T\textit{-modules}) \end{array}$$

Let B be the Brauer star algebra associated to the Brauer star with e edges and multiplicity 1. We have a surjection using the analogous mapping on respective sets:

$$\text{2-tilt}(B) \to \{\textit{simple-minded systems of } B\}.$$

Moreover, these maps are compatible with mutation on respective sets.

This article is structured as follows. In Section 2, we go through the basics of Brauer trees and Brauer tree algebras. We will collect essential results from [GR], [Rie2] and [CKL] about simple-minded systems of Brauer tree algebras that will be useful to us; and use these to indicate how the main theorem can be used to answer the Problem 1.2. In Section 3, we motivate the study of two-term tilting complex, and collect the facts and algorithms from [RS] and [SZI] about pointed Brauer tree and its relation with two-term tilting complexes of Brauer star algebra. In Section 4, we review various mutation theories, namely those of tilting complexes from [AI], those of simple-minded collection from [KY], those of Brauer tree from [Aih1], and those of simple-minded system from [Dug2] and [CKL]. We will present some propositions on these mutation theories in the case of Brauer tree algebras. Finally we prove Theorem 1.3 in Section 5.

Throughout this article, all the algebras are assumed to be finite dimensional over algebraically closed field, indecomposable, non-simple, and basic. In particular, they can be presented by quivers and relation. We work with right modules and write morphisms on the left, composing them from right to left. Likewise, paths in a quiver Q will be composed from right to left, and we often identify them with morphisms between projective right modules over (a quotient of) the path algebra. We will also assume the reader is familiar with the basics of Auslander-Reiten theory, and with Riedtmann's structuresatz [Rie1].

2. Brauer tree algebras and their simple-minded systems

Definition 2.1 (see for example [GR])**.** *A* Brauer graph G *is a finite undirected connected graph (possibly with loops and multiple edges) with the following data. To each vertex we assign a cyclic ordering of edges incident to it, and a positive integer called* multiplicity*.*

A Brauer tree *is a Brauer graph which is a tree, having at most one vertex with multiplicity greater than one. If there is such vertex, it is called* exceptional vertex*, otherwise we say the Brauer tree has trivial multiplicity. Traditionally, we choose the counter-clockwise direction as the cyclic ordering of edges; and denote the Brauer tree as* (G, v, m) *for a tree* G *with exceptional multiplicity* m *at the exceptional vertex* v*. For simplicity, we usually just use* G *as the notation for this triple. We will use* e *to denote the number of edges of* G*.*

A finite dimensional algebra A *is a* Brauer tree algebra *associated to a given Brauer tree* (G, v, m)*, if there is a one-to-one correspondence between the edges* j *of* G *and the simple* A*-modules* S_j *in such a way that the projective cover* P_j *of* S_j *has the following description. We have* $P_j/\operatorname{rad} P_j \cong \operatorname{soc} P_j \cong S_j$*, and the heart* $\operatorname{rad} P_j/\operatorname{soc} P_j$ *is a direct sum of two (possibly zero) uniserial modules* U_j *and* W_j *corresponding to the two vertices* u *and* w *at the end of the edge* j*. If the edges around* u *are cyclically ordered* $j, j_1, j_2, \dots, j_r, j$ *and the multiplicity of the vertex* u *is* m_u*, then the corresponding uniserial module* U_j *has composition factors (from the top)* $S_{j_1}, S_{j_2}, \dots, S_{j_r}, S_j, S_{j_1}, \dots, S_{j_r}, S_j, \dots, S_{j_r}$ *so that* $S_{j_1}, \dots, S_{j_r}$ *appear* m_u *times and* S_j *appears* $m_u - 1$ *times. We denote the basic algebra associated to a Brauer tree* G *as* $B^G_{e,m}$*.*

A Brauer star, which we usually denote by $\star$, is a Brauer tree where the underlying graph is a star, with exceptional vertex at the centre, the corresponding Brauer tree algebra is called Brauer star algebra. Note that the class of Brauer star algebras coincides with the class of symmetric Nakayama algebras.

For any $n \in \mathbb{Z}$, denote $\overline{n}$ to be the positive integer in $\{1, \dots, e\}$ with $n \equiv \overline{n} \mod e$. We fix a quiver and relation presentation for Brauer star algebra $B^\star_{e,m}$ associated to Brauer star with e edges and multiplicity m:

$$\begin{tikzcd}
1 \arrow[r, "\alpha_1"] & e \arrow[d, "\alpha_e"] \\
2 \arrow[u, "\alpha_2"] \arrow[r, dotted, no head] & \cdot
\end{tikzcd}$$

with relations given by setting all paths of length $em+1$ as zero. In particular, the radical of projective indecomposable P_i has projective cover $P_{\overline{i+1}}$. All indecomposable A-modules are uniserial, hence uniquely determined by its socle and Loewy length, and so we denote $M_{i,\ell}$ with $i \in \{1, \dots, e\}$ and $1 \leq \ell \leq em + 1$. For any i and any $\ell \leq em$, the Auslander-Reiten translate τ (which is isomorphic to the second syzygy functor Ω^2) sends $M_{i,\ell}$ to $M_{\overline{i+1},\ell}$. The Heller translate Ω, which is the inverse of suspension functor in the triangulated category $\underline{\text{mod}}$-A, sends $M_{i,\ell}$ to $M_{\overline{i-\ell+1},em+1-\ell}$; and inverse Heller transalte Ω^{-1} sends $M_{i,\ell}$ to $M_{\overline{i-\ell},em+1-\ell}$.

In [GR], Gabriel and Riedtmann classified the class of algebras that are stably equivalence to Brauer star algebra with e edges and multiplicity m, and this class is precisely the Brauer tree algebras which has the same number of edges e and same multiplicity m; the stable Auslander-Reiten (AR) quiver of this class of algebra is $\mathbb{Z}A_{em}/\langle\tau^e\rangle$. Riedtmann further investigated the stable equivalence between these algebras in [Rie2] and produced a description of the image of simple modules via a combinatorial gadget called configuration. To keep this article as concise as possible, by configuration we always mean Riedtmann's combinatorial configuration, which is equivalent to the original definition of configuration, see [CKL] for details.

Definition 2.2 ([Rie2]). *Let Δ be a stable representable quiver, a (combinatorial) configuration $\mathcal{C}$ is a set of vertices of Δ which satisfy the following conditions:*

1. *For any $e, f \in \mathcal{C}$,* $\mathrm{Hom}_{k(\Delta)}(e,f) = \begin{cases} 0 & (e \neq f), \\ k & (e = f). \end{cases}$
2. *For any $e \in \Delta$, there exists some $f \in \mathcal{C}$ such that* $\mathrm{Hom}_{k(\Delta)}(e,f) \neq 0$.

Remark 1. $k(\Delta)$ above is the mesh category of Δ. In the case of Brauer tree algebra, $\Delta = \mathbb{Z}A_{em}/\langle\tau^e\rangle$, regardless of the underlying Brauer tree. Therefore, for a Brauer tree algebra $B^G_{e,m}$ for some G, we say that a set $\mathcal{C}$ of vertices in $\mathbb{Z}A_{em}/\langle\tau^e\rangle$ is a *configuration of* $B^G_{e,m}$.

The motivation of this article comes from understanding simple-minded systems introduced by Koenig and Liu in [KL]. We denote by mod-A the category of all finitely generated right A-modules, by $\mathrm{mod}_{\mathcal{P}}A$ the full subcategory of mod-A whose objects have no nonzero projective direct summand, and by $\underline{\mathrm{mod}}$-A the stable category of mod-A modulo projective modules. Let $\mathcal{S}$ be a class of A-modules. The full subcategory $\langle\mathcal{S}\rangle$ of mod$A$ is the additive closure of $\mathcal{S}$. Denote $\langle\mathcal{S}\rangle * \langle\mathcal{S}'\rangle$ as the class of indecomposable A-modules Y such that there is a short exact sequence $0 \to X \to Y \oplus P \to Z \to 0$ with $X \in \langle\mathcal{S}\rangle$, $Z \in \langle\mathcal{S}'\rangle$, and P projective. Define $\langle\mathcal{S}\rangle_1 := \langle\mathcal{S}\rangle$ and $\langle\mathcal{S}\rangle_n := \langle\mathcal{S}\rangle_{n-1} * \langle\mathcal{S}\rangle$ for $n > 1$.

Definition 2.3 ([KL]). *Let A be self-injective algebra over a field. A class of objects $\mathcal{S}$ in $\mathrm{mod}_{\mathcal{P}}A$ is called a simple-minded system over A if the following conditions are satisfied:*

1. *(orthogonality condition) For any $S, T \in \mathcal{S}$,* $\underline{Hom}_A(S,T) = \begin{cases} 0 & (S \neq T), \\ \textit{division ring} & (S = T). \end{cases}$
2. *(generating condition) For all indecomposable non-projective A-module X, there exists some natural number n (depending on X) such that $X \in \langle\mathcal{S}\rangle_n$.*

In case A is representation-finite, we showed in [CKL] that every simple-minded system of A can be described by Riedtmann's configurations; we refer the reader to [CKL] for detailed study and survey on simple-minded systems for representation-finite self-injective algebras in general. We also recall from [CKL] that a class $\mathcal{S}$ of indecomposable A-modules is called a *simple-image* if there is a stable equivalence $\phi : \underline{\mathrm{mod}}\text{-}B \to \underline{\mathrm{mod}}\text{-}A$ such that $\mathcal{S}$ is the image of simple B-module under ϕ. If the stably equivalent algebra B is known, we say $\mathcal{S}$ is a *B-simple-image*. Obviously, a simple-image is always a simple-minded system. We extract the results from [CKL] and [Rie2] required in this article below.

Theorem 2.4. *Let $A = B^G_{e,m}$ be a Brauer tree algebra over an algebraically closed field k, and ${}_s\Gamma_A = \mathbb{Z}Q/\Pi$ be the stable AR quiver of A.*

1. *[Rie2] Any configuration $\mathcal{C}$ of A can be identified with a simple-image.*
2. *[CKL] There is a bijection:*

$$\begin{array}{rcl} \{\Pi\text{-}\textit{stable configurations of } \mathbb{Z}Q\} & \leftrightarrow & \{\textit{simple-minded systems of } A\} \\ \mathcal{C} & \mapsto & H(\mathcal{C}) \end{array}$$

for some map $H : k({}_s\Gamma_A) \to \underline{\mathrm{mod}}\text{-}A$ defined on the objects of the respective categories. Moreover, every simple-minded system of A appear as the restriction of image of the set of simple modules under some derived equivalence onto the stable module category of A.

(3) [CKL, Rie2] There is a bijection between the set of isomorphism classes of algebras stably equivalent to $B^{\star}_{e,m}$, and the set of orbits of simple-minded system under the action of stable auto-equivalences $\langle \Omega^n | n \in \mathbb{Z} \rangle$. In particular, any simple-minded system $\mathcal{S}$ of Brauer tree algebra, has its Ω-orbit splitting into at most two τ-orbits:

$$\{\Omega^n \mathcal{S} | n \in \mathbb{Z}\} = \{\tau^n \mathcal{S} | n \in \mathbb{Z}\} \sqcup \{\tau^n \Omega^{-1} \mathcal{S} | n \in \mathbb{Z}\}$$

Corollary 2.5. *Let A be a Brauer star algebra and $F : D^b(\mathrm{mod}\text{-}A) \to D^b(\mathrm{mod}\text{-}B)$ be a derived equivalence. Then there is a two-term tilting complex T concentrated in degree 0 and -1 which gives rise to the following diagram.*

$$\begin{tikzcd}
\underline{\mathrm{mod}}\text{-}E_T \arrow[r, "\overline{F_T}"] & \underline{\mathrm{mod}}\text{-}A & \underline{\mathrm{mod}}\text{-}B \arrow[l, "\overline{F}"'] \\
\{\textit{simple } E_T\textit{-modules}\} \arrow[r, mapsto] & \mathcal{S} & \{\textit{simple } B\textit{-modules}\} \arrow[l, mapsto]
\end{tikzcd}$$

where $\overline{F_T}$ and $\overline{F}$ are the induced stable equivalences and $n \in \mathbb{Z}$. In particular, Problem 1.2 has a positive answer when A is a Brauer star algebra.

Proof. This follows from combining Theorem 1.3 and Theorem 2.4. □

In the remaining of this article, we fix the following coordination for the stable AR-quiver of Brauer tree algebras using the pair appearing in the subscript of an indecomposable $B^{\star}_{e,m}$-module. Note that the stable AR-quiver of Brauer tree algebra $B^{G}_{e,m}$ is isomorphic to the stable tube $\mathbb{Z}A_{em}/\langle \tau^e \rangle$. The simple $B^{\star}_{e,m}$-modules lie on the bottom rim of the stable AR-quiver, and radical of projective indecomposable $B^{\star}_{e,m}$-modules lie on the top rim of the stable AR-quiver.

Example 2. The following is the stable AR-quiver of $B^{\star}_{3,2}$ (we omit the symbol M):

$$\begin{tikzcd}[column sep=tiny]
 & & & & & (3,6) \arrow[dr] & & (2,6) \arrow[dr] & & (1,6) \arrow[dr] & & (3,6) \\
 & & & & (3,5) \arrow[ur] \arrow[dr] & & (2,5) \arrow[ur] \arrow[dr] & & (1,5) \arrow[ur] \arrow[dr] & & (3,5) \arrow[ur] & \\
 & & & (3,4) \arrow[ur] \arrow[dr] & & (2,4) \arrow[ur] \arrow[dr] & & (1,4) \arrow[ur] \arrow[dr] & & (3,4) \arrow[ur] & & \\
 & & (3,3) \arrow[ur] \arrow[dr] & & (2,3) \arrow[ur] \arrow[dr] & & (1,3) \arrow[ur] \arrow[dr] & & (3,3) \arrow[ur] & & & \\
 & (3,2) \arrow[ur] \arrow[dr] & & (2,2) \arrow[ur] \arrow[dr] & & (1,2) \arrow[ur] \arrow[dr] & & (3,2) \arrow[ur] & & & & \\
(3,1) \arrow[ur] & & (2,1) \arrow[ur] & & (1,1) \arrow[ur] & & (3,1) \arrow[ur] & & & & &
\end{tikzcd}$$

$\{(1,1),(2,1),(3,1)\}$ is the set of simple $B^{\star}_{3,2}$-module. Another example of simple-minded system (i.e. configuration) is $\mathcal{S} = \{(1,1),(2,3),(3,5)\}$, the (unique) Brauer tree algebra B such that $\mathcal{S}$ is a B-simple-image is the one associated to the graph (tree) of a line with exceptional vertex at the end of the line, i.e. $\bullet \text{---} \circ \text{---} \circ \text{---} \circ$.

We will use this coordination throughout the article. Note that our coordination is slightly different from the conventional one, where the "x-axis" goes from left to right, the transformation from our coordination to the conventional one is $(x,y) \mapsto (em - x, y)$.

Theorem 2.4 says that in order to understand simple-minded systems over representation-finite self-injective algebras, we can study configurations instead, transforming most of the algebraic exposition to a combinatorial one. In the following, we play with this combinatorics to give us some tools which will be useful later in the proof of the main theorem. An *extremal vertex* of Brauer tree G is a vertex of valency 1; we call the edge connected to an extremal vertex as *leaf.*

Lemma 2.6. *Let G be a Brauer tree and S a simple $B^G_{e,m}$-module. Then S lies on the rim of the stable AR-quiver, if and only if, the edge in G which corresponds to S is a leaf attached to a non-exceptional extremal vertex.*

Proof. By the construction of Brauer tree algebras, an edge is a leaf attached to a non-exceptional extremal vertex if and only if the corresponding indecomposable projective module is uniserial. Also note that for any simple $B^G_{e,m}$-module S, whose projective cover is P, we have almost split sequence starting at $\Omega(S)$ being:

$$0 \to \Omega(S) \to P \oplus \operatorname{rad}(P)/\operatorname{soc}(P) \to \Omega^{-1}(S) \to 0$$

This says that P is uniserial if and only if, $\Omega(S)$ and $\Omega^{-1}(S)$ is on a rim of the stable AR-quiver, say located (without loss of generality) at $(1, em), (e, em)$ respectively, which in turns is equivalent to S located at $(1, 1)$, i.e. another rim of the stable AR-quiver. □

We need some combinatorial lemma about configurations:

Lemma 2.7 ([Rie3], Lemma 2.5)**.** *A set $\mathcal{C}$ in the vertex set $(\mathbb{Z}A_n)_0$ of $\mathbb{Z}A_n$ is a configuration of $\mathbb{Z}A_n$ if and only if*

$$\omega_n\mathcal{C} \cup \tau^{(n+1)\mathbb{Z}}(0, 1)$$

is a configuration of $\mathbb{Z}A_{n+1}$, where ω_n is an injection $\omega_n : (\mathbb{Z}A_n)_0 \to (\mathbb{Z}A_{n+1})_0$ given by:

$$\omega_n(x, y) = \begin{cases} (x, y) & \text{if } 0 \le x - y < x \le n \\ (x, y + 1) & \text{if } 0 < x < y \end{cases}$$

and by the rule $\omega_n\tau^n = \tau^{n+1}\omega_n$.

Applying this lemma on the class of Brauer tree algebras, i.e. for configurations of $\mathbb{Z}A_{em}$ which are stable under $\tau^{e\mathbb{Z}}$, one can see the lemma can be refined to:

Lemma 2.8. *A set $\mathcal{C}$ in $(\mathbb{Z}A_{em}/\langle\tau^e\rangle)_0$ is a configuration of Brauer tree algebra associated to Brauer star algebra $B^\star_{e,m}$ if and only if*

$$\mathcal{C}^+ := \omega^{(m)}_e\mathcal{C} \cup \{(e + 1, 1)\}$$

is a configuration of $B^\star_{e-1,m}$ where $\omega^{(m)}_e$ is an injection $\omega^{(m)}_e : (\mathbb{Z}A_{em})_0 \to (\mathbb{Z}A_{(e+1)m})_0$ given by:

$$\omega^{(m)}_e(x, y) = (x, y + \ell)$$

when $-\ell e \le x - y < (1 - \ell)e$ for some $\ell \in \{0, \dots, m\}$.

When we compare two simple-minded systems, both of which appear to be B-simple-image for some algebra B, we want to be able to tell whether they lie in the same τ-orbit (c.f. Theorem 2.4). For this purpose, we need to refine the above combinatorial lemma.

Suppose we have a configuration $\mathcal{C}$ of Brauer star algebra $B^\star_{e,m}$ which corresponds (under the map of Theorem 2.4) to a $B^G_{e,m}$-simple-image, each $(x, y) \in \mathcal{C}$ lying on the rim of stable AR-quiver (i.e. (x, y) with $y = 1$ or em) then corresponds to a leaf of G by Lemma 2.6. Fix any one of such (x, y), let G' be the Brauer tree obtained by removing a leaf corresponding to (x, y) in G. Apply Heller shifts Ω^n (for some n) to $\mathcal{C}$ so that $(x, y) \mapsto (1, 1)$. More explicitly, apply τ^{-x+1} (i.e. $n = -2x + 2$) if $y = 1$, or $\tau^{-x+1}\Omega^{-1}$ (i.e. $n=-2x+1$) if $y=em$. Now we have $\Omega^n\mathcal{C}=\mathcal{D}^+$ for some configuration $\mathcal{D}$ which is a $B^{G'}_{e-1,m}$-simple-image. Obviously, $\Omega^{-n}\mathcal{D}$ is still a configuration of $B^{G'}_{e-1,m}$. We call this process as "*cutting off a leaf*".

The following observation is crucial for proving the main theorem:

Corollary 2.9 (Tree Pruning Lemma)**.** *Suppose G is a Brauer tree with e edges and multiplicity m, where the valency of the exceptional vertex is ℓ. Let $\mathcal{C}$ be the configuration of $B^G_{e,m}$ representing the simple $B^G_{e,m}$-module. Then the effect on $\mathcal{C}$ after successively cutting off leaves of G until reaching $B^\star_{\ell,m}$ is either $\mathcal{C}^\star_{\ell,m}:=\{(1,1),\dots,(\ell,1)\}$ or $\Omega^{-1}\mathcal{C}^\star_{\ell,m}=\{(1,\ell m),\dots,(\ell,\ell m)\}$, depending only on the τ-orbit for which $\mathcal{C}$ lie in.*

By the virtue of this result, we can classify configurations of Brauer tree algebras into two types. A configuration $\mathcal{C}$ which is a $B^G_{e,m}$-simple-image is said to be of "bottom-type" (resp. "top-type") if the resulting configuration after tree pruning is $\mathcal{C}^\star_{\ell,m}$ (resp. $\Omega^{-1}\mathcal{C}^\star_{\ell,m}$).

3. Tilting complexes and Pointed Brauer tree

The other structure we are interested in is the (one-sided) tilting complexes of Brauer star algebra $B^\star_{e,m}$. Let us first recall some necessary definitions. The following "definition" of tilting complex is given by results of [AI] and [KY], equivalent to the original one given by Rickard [Ric2].

Definition 3.1. *Let A be a finite dimensional self-injective algebra. A (one-sided)* tilting complex *is an object T in the bounded homotopy category $\mathcal{T}=K^b(\text{proj-}A)$ of finitely generated projective A-modules if*

1. $\mathrm{Hom}_{\mathcal{T}}(T,T[i])=0$ *for all* $i\neq 0$,
2. $\mathcal{T}=\mathrm{thick}(T)$,
3. *The indecomposable summands of T are pairwise non-isomorphic and the number of indecomposable summands of T equals to the rank of Grothendieck group of $\mathcal{T}$.*

An object in $\mathcal{T}$ is a two-term *complex if it is concentrated in at most two consecutive homological degree. A tilting complex is* two-restricted *if it is a direct sum of indecomposable two-term complexes.*

Given a tilting complex T, let E_T denote the derived equivalent algebra $\mathrm{End}_{\mathcal{T}}(T)$, and F_T be the triangulated equivalence $\mathrm{Hom}_{\mathcal{T}}(T,-)$ or the derived equivalence $\mathrm{Hom}_{D^b(\text{mod-}A)}(T,-)$. By a result of Rickard [Ric1], any algebra derived equivalent to a Brauer tree algebra is also a Brauer tree algebra, hence, we sometimes call a tilting complex to be "star-to-tree tilting complex" if A is the Brauer star algebra $B^\star_{e,m}$. However, there is an abundance of tilting complexes (even up to shifts and homotopy equivalences), and we should restrict to a much more refined subclass when studying the homological theories around these algebras. Our choice in the current article is the set of two-term tilting complexes. The reason comes from the work of Abe and Hoshino [AH], where they showed that every derived equivalence between self-injective algebras given by a tilting complex is a composition of derived equivalences given by two-term tiltings. Moreover, in [AIR] it is shown that the set of two-term tilting complexes of symmetric algebra is in bijection with the set of functorially finite torsion classes of its module category, emphasising the importance of two-term tilting complexes in the study of homological behaviour of symmetric algebra. We also remark that these statements are also true for any finite dimensional self-injective algebras by inserting "Nakayama-stable" into appropriate places, c.f. [AH] and [Miz].

Since Brauer tree algebras are encoded by the rich combinatorics in the Brauer trees, it is natural to seek for combinatorial construction and description of two-term tilting complexes. One approach is given as follows by Schaps and Zakay-Illouz. Some other approaches to this include a combinatorial one from [AZ], and using a recently invented theory of support τ-tilting module [Ada] along with the bijection between support τ-tilting modules and two-term tilting complexes [AIR]. We will only use Schaps-Zakay-Illouz approach in this article. We first recall the following combinatorial gadget:

Definition 3.2 ([SZI]). *Let (G, v, m) be a Brauer tree with e edges. A* pointing *is a set of data assigned to each non-exceptional vertex u, of a pair of edges i, j which are adjacent in the cyclic ordering at u (i.e. a* section *around u). We call this pair of edges as* pointed section*. If u is extremal attached to leaf i, then we take (i, i) as the required pair. The tree G together with a pointing is called a* pointed Brauer tree*.*

We recall the main result of Schaps and Zakay-Illouz [SZI]:

Theorem 3.3 (Schaps-Zakay-Illouz correspondence [SZI]). *There is a one-to-one correspondence between the set of two-restricted tilting complexes (up to shifts and homotopy equivalences) for the Brauer star algebra $B^{\star}_{e,m}$ associated to a Brauer star with e edges and multiplicity m, and the set of pointed Brauer trees having e edges and multiplicity m along with a marked non-exceptional vertex. Moreover, under this correspondence, the underlying Brauer tree of the Brauer tree algebra E_T coincide with the underlying Brauer tree of the pointed Brauer tree corresponding to T.*

Remark 3. From the definition of pointing, there is only one pointing on the Brauer star, this corresponds to the algebra $B^{\star}_{e,m}$ itself considered as a tilting complex.

Looking closely at the combinatorics and the bijection, [RS] gave a description of two-term star-to-tree tilting complexes (those they termed "completely folded") using *alternating pointing.* For each Brauer tree, apart from the star, with a marked non-exceptional vertex, there are precisely two kinds of alternating pointing. Let us be more precise with the construction of such pointed Brauer trees and their associated tilting complexes.

3.1. The Green's walk around Brauer tree.

Let us first very quickly review the Green's walk [Gre]. As we have chosen the counter-clockwise direction for the cyclic ordering of the edges, the Green's walk around G is "a counter-clockwise walk along the edges of G". We start by picking a vertex with an adjacent edge and run the following recursion step until going back to the starting point (same vertex same edge). The recursion step can be stated simply as: Given a vertex u and edge i, one obtains a new vertex v as the other end of i and a new edge j as the next edge at v in the cyclic ordering. When one assigned a pointing on G, we say a stage, e.g. (u, i), of the Green's walk have *reached a pointed section* if the pointing at u is (j, i).

3.2. Constructing alternating pointing.

For any vertex $u \in G$, recall the distance $d(u)$ is the minimum number of edges from the exceptional vertex to u. Let G be the Brauer tree, and start at an edge sharing the exceptional vertex. The following algorithm constructs one kind of alternating pointing. The algorithm iterates using the Green's walk, starting from the pair (u, i) with i be an edge attached to the exceptional vertex:

(1) Let $i, i_1, \ldots, i_\ell$ be the cyclic ordering of the edge incident to the vertex u. If $d(u)$ odd (resp. even), then assign pointing (i, i_1) (resp. (i_ℓ, i)) at u, we say this is "pointing in the first (resp. last) section at u".

(2) Go around the tree using the Green's walk, until reaching the next non-exceptional vertex u' which is not assigned a pointing.

(3) Repeat Step (1) by replacing u with u', until all vertices are assigned with a pointing.

For convenience, we call this alternating pointing as *odd-first pointing*. The other alternating pointing is given by pointing in first (resp. last) section when $d(u)$ even (resp. odd), hence we call it *even-first pointing*. Also note that if the vertex is a leaf of a tree, then the pointing is unique.

3.3. **Constructing two-term tilting complexes.** Now given an alternating pointed Brauer tree with a marked non-exceptional vertex v, we number the non-exceptional vertices according to the following iteration using the Green's walk, starting at the pointed section at (v, i), where $v = u$ and i is the edge attached to vertices v and v', with v being furtherer away from the exceptional vertex, and $n = 1$.

(1) If we reach a pointed section at u, assign the number n to u and then replace n with $n + 1$. Otherwise, do nothing.
(2) Go around the tree using Green's walk, until reaching the next vertex u' which is not assigned a number.
(3) Repeat Step (1) by replacing u with u', until all non-exceptional vertices are assigned with a number.

Now label each edge by the same number as the vertex farthest from the exceptional vertex. Define a (two-term) star-to-tree tilting complex $T = \bigoplus_{i=1}^{e} T_i \in$ 2-tilt$(B^\star_{e,m})$ by the following recursion on the distance from exceptional vertex.

(1) If edge i is adjacent to the exceptional vertex, then T_i is the stalk complex $0 \to P_i \to 0$ with P_i at degree 0 (resp. -1) if pointing is odd-first (resp. even-first).
(2) Let $i_1, \ldots, i_\ell$ be the edges adjacent to the chosen vertex u, with i_1 be an edge with T_{i_1} assigned. For each $j \neq 1$, we have two cases:
    (a) If P_{i_1} is in homological degree 0, then assign T_{i_j} with the two-term complex $P_{i_j} \to P_{i_1}$ concentrated in degree -1 and 0.
    (b) If P_{i_1} is in homological degree -1, then assign T_{i_j} with the two-term complex $P_{i_1} \to P_{i_j}$ concentrated in degree -1 and 0.

For convenience in later part of the article, if the pointing is even-first, then replace the corresponding tilting complex $T \in$ 2-tilt$(B^\star_{e,m})$ by its shift $T[-1]$. With this choice of homological positions, the homological degree 0 component of all the indecomposable summands in T will always be non-zero. We say that a two-term star-to-tree tilting complex (excluding $B^\star_{e,m}$ itself) appearing in this form, which corresponds to a even-first (resp. odd-first) pointed Brauer tree, is *positive* (resp. *negative*), in order to emphasies the degrees where non-zero modules appear. Consequently,

$$\text{2-tilt}(B^\star_{e,m}) = \{\text{negative two-term tilting}\} \cup \{T[1] | T \text{ positive two-term tilting}\} \cup \{B^\star_{e,m}, B^\star_{e,m}[1]\}.$$

Example 4. Consider the following pointed Brauer tree with the odd-first pointing, where the exceptional vertex is coloured black. Following [SZI], the pointing at each vertex is labelled by a black dot in the pointed section. Note that at the leaves of the Brauer tree, the pointing is unique so we do not need to put any black dot. We choose the marked vertex as the one of valency 3 here. For clarity, we draw the Green's walk around the tree with the dotted line in Figure 1, starting at the pointed section of the marked vertex.



FIGURE 1. Pointed Brauer tree and a Green walk.



FIGURE 2. Labelling of vertices and edges on a pointed Brauer tree.

The labelling on the vertices and on the edges are shown in the figure 2. The Schapes-Zakay-Illouz correspondence gives the following two-term tilting complex:

$$\begin{array}{rcl} 0 & \to & P_2 \\ \oplus(P_3 & \to & P_2) \\ \oplus(\ \ 0 & \to & P_4) \\ \oplus(P_1 & \to & P_4) \\ \oplus(P_1 & \to & P_5) \\ \oplus(P_1 & \to & P_6) \end{array}$$

with indecomposable projective $B^{\star}_{6,m}$-modules P_2, P_4, P_5, P_6 appearing in homological degree 0, and all other indecompoable projecitve modules appearing in homological degree -1.

We end this section by answering Problem 1.1 for Brauer star algebras.

Theorem 3.4. *Suppose A is an algebra derived equivalent to $B^{\star}_{e,m}$. There is a two-term tilting complex T concentrated in degree 0 and -1, so that E_T is Morita equivalence to A. In particular, Problem 1.1 has a positive answer for Brauer star algebras.*

Proof. It is well-known that any such A is a Brauer tree algebra associated to Brauer tree G with e edges and multiplicity m. Now we can pick any alternating pointing and any non-exceptional vertex of G, and construct two-term tilting complex T, which has E_T Morita equivalent to A. The statement now follows from Theorem 3.3. $\square$

4. MUTATION THEORIES

One approach to construct tilting complexes is via iterative mutations starting from A. One can also construct simple-minded collections and simple-minded systems similarly using mutation starting from the set of simple modules. We now go through the definitions and lemmas regarding mutations of these objects. Note that we use the term tilting mutation instead of silting mutation in [AI] as all silting complexes are tilting complexes for a weakly symmetric algebra A (such as Brauer tree algebras). Throughout this section the algebras we consider are always (weakly) symmetric. Following the convention in [CKL], we use Roman fonts for tilting complexes, bold fonts for simple-minded collections, and calligraphic fonts for simple-minded systems.

4.1. Mutation of tilting complexes and simple-minded collections.

Definition 4.1 (Mutation of tilting complex [AI]). *Let* $T = T_1 \oplus \cdots \oplus T_r$ *be a tilting complex and* $T = X \oplus M$*, a left tilting mutation of* T *at* X*, denoted* $\mu_X^-(T) := T_1' \oplus \cdots \oplus T_r'$ *is given by*

1. $T_i' = T_i$ *if* T_i *a summand of* X*.*
2. *Otherwise,*
$$T_i' := \mathrm{cone}(\textit{minimal left } \mathrm{add}(M)\textit{-approximation of } T_i)$$

A right tilting mutation μ_X^+ *is defined similarly using right approximation. A tilting mutation is said to be* irreducible *if* X *is indecomposable.*

Some of the results in this section work for both left and right mutations, in which case we simplify our notation for mutations by superscripting with ±. We note the following useful fact, which says that left and right mutations are operations inverse to each other:

Proposition 4.2 ([AI], Prop 2.33). *Suppose* T *is a tilting complex and* $T = X \oplus M$*, writing* $\mu_X^+(T) = Y \oplus M$ *and* $\mu_X^-(T) = Z \oplus M$*, then we have* $\mu_Y^-\mu_X^+(T) = T = \mu_Z^+\mu_X^-(T)$*.*

By construction, tilting mutation respects derived equivalence as follows. Given derived equivalence $F : D^b(\text{mod-}A) \to D^b(\text{mod-}B)$ (or triangulated equivalence $F : K^b(\text{proj-}A) \to K^b(\text{proj-}B)$, and a tilting complex $T \in K^b(\text{proj-}A)$ with a summand X of T, we have $F(\mu_X^\pm(T)) = \mu_{FX}^\pm(FT)$.

We will not work with simple-minded collections most of the time, but they serve as a bridge between tilting complexes and simple-minded systems, so we state the necessary definitions and results here, and refer the reader to [CKL] for more details.

Definition 4.3 (Simple-minded collection and its mutations [KY]). *Let* A *be a weakly symmetric algebra, and* $\mathbf{S} = \{X_1, \dots, X_r\}$ *a collection in the bounded derived category* $D^b(\text{mod-}A)$*.*

1. $\mathbf{S}$ *is a* simple-minded collection *if for all* $i, j = 1, \cdots, r$*, the following conditions are satisfied:*
    - (a) *(orthogonality)* $Hom(X_i, X_j) = k\delta_{i,j}$
    - (b) *(generating)* $D^b(\text{mod-}A) = thick(X_1 \oplus \cdots \oplus X_r)$*.*
    - (c) *(silting/tilting)* $Hom(X_i, X_j[m]) = 0$ *for any* $m < 0$*.*
2. *Let* $\mathbf{X} \subset \mathbf{S}$*,* a left mutation *of* $\mathbf{S}$ *at* $\mathbf{X}$*, denoted* $\mu_{\mathbf{X}}^-(\mathbf{S}) := \{Y_1, \dots, Y_r\}$ *is given by*
    - (a) $Y_i = X_i[1]$ *if* $X_i \in \mathbf{X}$*.*
    - (b) *Otherwise,* Y_i *is the cone of the minimal left* $\mathcal{F}(\mathbf{X})$*-approximation of* $X_i[-1]$*, where* $\mathcal{F}(\mathbf{X})$ *is the extension closure of* $\mathbf{X}$*.*

    A right tilting mutation $\mu_{\mathbf{X}}^+$ *is defined similarly using right approximation. A mutation of simple-minded collection is said to be* irreducible *if* $\mathbf{X}$ *has cardinality 1.*

The mutation of simple-minded collection also respects derived equivalence, namely $F(\mu_{\mathbf{X}}^\pm(\mathbf{S})) = \mu_{F\mathbf{X}}^\pm(F\mathbf{S})$ for derived equivalence $F : D^b(\text{mod-}A) \to D^b(\text{mod-}B)$.

One of the main results in Al-Nofayee [AN] is that for a Nakayama-stable simple-minded collection $\mathbf{S}$, one can construct a tilting complex T so that $F_T(\mathbf{S})$ is the set of simple E_T-modules. This is also shown in [KY]. The key result from [KY] is the following.

Theorem 4.4 (Koenig-Yang bijection [KY]). *Suppose* A *is a finite dimensional symmetric algebras. There is a one-to-one correspondence between the equivalence classes of tilting objects in* $K^b(\text{proj-}A)$ *and the equivalence classes of simple-minded collections in* $D^b(\text{mod-}A)$*. Moreover, this bijection respect mutations on respective classes.*

Remark 5. Two tilting objects are equivalence if they are homotopy equivalent up to shifts. Two simple-minded collections are equivalence if their extension closure coincides.

Even more specifically, let T be tilting complex corresponding to simple-minded collection $\mathbf{S}$, then there is an ordering on indecomposable summands of T and elements of $\mathbf{S}$ such that for any subset $\mathbf{X} = \{X_{i_1}, \dots, X_{i_\ell}\} \subset \mathbf{S}$ and summand $X = T_{i_1} \oplus \cdots \oplus T_{i_\ell}$, $\mu^{\pm}_{\mathbf{X}}(\mathbf{S})$ is in correspondence with $\mu^{\pm}_{X}(T)$. Note that under the Koenig-Yang bijection, the set of simple A-modules (considered as a set of stalk complexes concentrated in homological degree 0) corresponds to the canonical tilting complex A.

In [AI], Aihara and Iyama posed a question asking whether one can obtain all the tilting complexes just by iterative irreducible mutations (or by iterative irreducible left mutations) starting from the canonical tilting complex A. Aihara affirmed this for representation-finite A:

Theorem 4.5 ([Aih2]). *Let A be representation-finite symmetric algebra and T be a tilting complex concentrated in non-positive homological degrees, then T can be obtained by iterative irreducible left mutations, i.e. $T = \mu^-_{M_\ell}\mu^-_{M_{\ell-1}} \cdots \mu^-_{M_1}(A)$ for some indecomposable partial tilting complexes $M_1, \dots, M_\ell$.*

Remark 6.
1. Any tilting complex can be shifted so that it is concentrated in non-positive homological degree.
2. By Koenig-Yang bijection, this is equivalent to saying that any Nakayama-stable simple-minded collection (possibly after shifting) can be obtained by iterative left mutation starting from the set of simple A-modules.
3. Dually, if T is concentrated in non-negative homological degrees, then T can be obtained by iterative irreducible right mutations with $T = \mu^+_{N_\ell} \cdots \mu^+_{N_1}(A)$.

4.2. **Mutation of Brauer trees.** Mutation of tilting complex can be reformulated as mutation on the class of derived equivalent algebras: for a summand P of an algebra A, let T be the tilting mutation $\mu^-_X(A)$, then we can define the left algebra mutation as the algebra E_T. On the class of Brauer tree algebra, this gives a mutation on the Brauer trees. Using the result from [Aih1, Theorem 1.5] or [Dug1, Section 2,3], it turns out we can define mutation of Brauer tree algebras directly from the underlying combinatorics.

Definition 4.6 (Mutation of Brauer tree). *Let G be a Brauer tree, and i be an edge of G, the* left *mutation of G at i, denoted $\mu^-_i(G)$, can be constructed as follows. Suppose the vertices attached to i are u and v, with j and k being the previous edges in the cyclic ordering around u and v respectively. The mutated tree is given by removing the edge i from G, and replace with an edge i' connected to j and k. In particular, if (without loss of generality) u is only of valency one (i.e. an extremal vertex), then i' is attached to u again.*

Similarly, define the right mutation $\mu^+_i(G)$ by removing i and connecting i' to the next *edges in the cyclic ordering around u and v. The two mutations can be visualised as in Figure 3.*

The discussion prior to this definition says that, if T is a star-to-tree tilting complex which takes the Brauer star to a Brauer tree G, and let i be an edge correspond to a summand X of T, then $\mu^{\pm}_X(T)$ is a star-to-tree tilting complex which takes the Brauer star to the mutated Brauer tree $\mu^{\pm}_i(G)$.

We now present an algorithm to mutate from the Brauer star to arbitrary Brauer tree so that the mutated tilting complex under the corresponding mutation is always two-term. We start with some sufficient criteria for a mutated tilting complex to be a two-term.

Lemma 4.7. *Let T be a tilting complex concentrated in non-positive homological degrees, and X be an indecomposable summand of T, we have the following:*



FIGURE 3. Mutation of Brauer tree.

(1) If $\mu_X^-(T)$ is two-term, then so is T.
(2) If T is two-term and M is a stalk complex $0 \to P_i \to 0$ for some $i \in \{1, \dots, e\}$ concentrated in homological degree 0, then $\mu_X^-(T)$ is two-term.

Proof. (1): Recall from [AI] that there is a partial order on the tilting complexes defined by $T \geq U$ if $\mathrm{Hom}_{\mathcal{T}}(T, U[i]) = 0$ for all $i > 0$. As T and $\mu_X^- N(T)$ are both concentrated in non-positive degrees, it follows from [Aih2, Proposition 2.9] that, $A \geq \mu_X^-(T) \geq A[1]$ and $A \geq T \geq A[\ell]$ for some $\ell \geq 1$. We also have $T \gneqq \mu_X^-(T)$ from [AI, Theorem 2.35]. These combine to give $A \geq T \gneqq \mu_X^-(T) \geq A[1]$, and so $\ell = 1$, which means T is two-term by [Aih2, Proposition 2.9].

(2): This is easy to see from the definition of mutation, for $T = X \oplus M$ with $\mu_X^-(T) = Y \oplus M$, then Y is the cone of a morphism from stalk complex concentrated in degree 0 to a two-term complex concentrated in degree 0 and -1, so Y is two-term as well. □

Remark 7. If T is concentrated in non-negative degrees instead, then the analogous results hold for right mutations instead of left mutations.

We will be implicitly using a correspondence between edges of a Brauer tree G and indecomposable summands of a star-to-G tilting complex T. More precisely, for E_T a Brauer tree algebra associated to G, and X an indecomposable summand of T, $F_T(X)$ is an indecomposable projective $B_{e,m}^G$-module, corresponding to an edge of G. We also note that, by Schape-Zakay-Illous correspondence, the each indecomposable stalk complex summand of a two-term tilting complex T always corresponds to an edge attached to the exceptional vertex.

Theorem 4.8. *Suppose T is a negative two-term tilting complex of $B_{e,m}^{\star}$ associated to Brauer tree G, where the valency of exceptional vertex of G is $e - \ell$, then T can be obtained by ℓ irreducible left mutations at stalk complexes. Dually, the same result is true if one replaces negative and left by positive and right respectively.*

Proof. By the discussion in the paragrah above and the previous lemma, it suffices to prove the following combinatorial result: Any Brauer tree G with $e - \ell$ valency at the exceptional vertex can be obtained by ℓ left mutations at edges attached to the exceptional vertex. Thanks to Proposition 4.2, this can be proved by finding an algorithm to obtain the Brauer star from the Brauer tree using right mutations, such that after each mutation, the valency of exceptional vertex is increased by 1.

Given an edge i of G connected to the exceptional vertex, we can define a branch of G connected to i as the following subtree of G. If u is the exceptional vertex itself, or u is a non-exceptional vertex for which there is a path from u to the exceptional vertex, ending at edge i, then u is in the branch of i. The edges of the branch connected to i are all the edges of G for which both ends are as described before. In particular, a branch is a Brauer tree where the exceptional vertex is of valency 1.

Our algorithm is to repeat the following recursively on every branch of the tree G: take a branch of the tree G connected to $i = i_0$ which contain more than one edge, let the nonexceptinal end of i_0 be u and suppose $i_0, i_1, \dots, i_n$ (note $n \geq 1$ always) is the cylic ordering of edges around u, we then right mutate at i_n. According to the mutation rule of Brauer trees, i_n is then removed and replaced by an edge with one ends connected to the exceptional vertex. Therefore the valency of exceptional vertex has increased precisely by 1. □

Example 8. In Figure 4, we show how the algorithm is applied to our previous example 4. Note



FIGURE 4.

that the Brauer star is not ordered in the presentation we fixed in the beginning. In order to get back the correct ordering, we apply the permutation $\pi = (156)$ to the labelling. Note that we have to fix the edges labelling on the edges attached to the exceptional vertex. This changes the mutation sequence to $\mu_5^- \mu_6^- \mu_1^- \mu_3^-$. Consequently, the two-term tilting complex T corresponding to this odd-first pointed Brauer tree, with marked vertex as the one of valency 3, is equal to the mutated complex: $\mu_{P_5}^- \mu_{P_6}^- \mu_{P_1}^- \mu_{P_3}^-(A)$.

4.3. Mutation of simple-minded systems.

Definition 4.9 (Mutation of simple-minded system [Dug2, CKL]). *Let $\mathcal{S} = \{X_1, \dots, X_r\}$ be a simple-minded system (of a weakly symmetric algebra) and $\mathcal{X} \subset \mathcal{S}$, a left mutation of $\mathcal{S}$ at $\mathcal{X}$, denoted $\mu_{\mathcal{X}}^-(\mathcal{S}) := \{Y_1, \dots, Y_r\}$ is given by*

1. *$Y_i = \Omega^{-1} X_i$ if $X_i \in \mathcal{X}$*
2. *Otherwise, Y_i is the cone of the minimal left $\mathcal{F}(\mathcal{X})$-approximation of ΩX_i, where $\mathcal{F}(\mathcal{X})$ is the extension closure of $\mathcal{X}$.*

A right mutation $\mu_{\mathcal{X}}^+$ is defined similarly using right approximation. A tilting mutation is said to be irreducible *if X is indecomposable.*

Remark 9. Again, mutation respects stable equivalence. Moreover, for any simple-minded system $\mathcal{S}$ of A and $\mathcal{X} \subset \mathcal{S}$, if we have derived equivalence $F : D^b(\text{mod-}A) \to D^b(\text{mod-}B)$, then we have $\overline{F}(\mu^{\pm}_{\mathcal{X}}(\mathcal{S})) = \mu^{\pm}_{\overline{F}\mathcal{X}}(\overline{F}\mathcal{S})$, where $\overline{F} : \underline{\text{mod}}\text{-}A \to \underline{\text{mod}}\text{-}B$ is the induced stable equivalence.

Example 10. Let A be the Brauer star algebra $B^{\star}_{2,m}$, and $\mathcal{S}_A = \{S_1, S_2\} = \{(1,1),(2,1)\}$ be the set of simple A-modules. Then $\mu^-_{S_1}(\mathcal{S}_A) = \{(2,2m),(1,2)\}$, which is in fact $B^G_{e,m}$-simple-image, where G is the Brauer line with one extremal vertex being exceptional.

Let $\eta_A : D^b(\text{mod-}A) \to \underline{\text{mod}}\text{-}A$ be the canonical quotient functor. Note that any (Nakayama-stable) simple-minded collection $\mathbf{S} = \{X_1, \ldots, X_r\}$ restricts to a simple-minded system in the stable module category: $\eta_A(\mathbf{S}) = \{\eta_A(X_1), \ldots, \eta_A(X_r)\}$. The following result from [Dug2] provides a connection between mutation of simple-minded systems and other structures.

Proposition 4.10 ([Dug2], Proposition 5.4)**.** *Let A be a self-injective algebra, $\mathbf{S} = \{S_1, \ldots, S_r\}$ be the set of simple A-modules, regarded as stalk complex concentrated in degree 0, and $\mathbf{X} = \{S_{i_1}, \ldots, S_{i_\ell}\}$ with $\nu\mathbf{X} = \mathbf{X}$ where ν is the Nakayama functor of A. Then*

$$\eta_A(\mu^{\pm}_{\mathbf{X}}(\mathbf{S})) = \mu^{\pm}_{\eta_A(\mathbf{X})}(\eta_A(\mathbf{S})). \tag{4.1}$$

Combining with Koenig-Yang bijection, and Remark 9, the formula (4.1) is true for any Nakayama-stable simple-minded collection. In another words, mutation allows one to construct simple-minded systems of A which appear to be a B-simple-image for every B derived equivalent to A via a tilting complex which can be obtained by iterative mutations. In [CKL], the author, along with Koenig and Liu, have shown that any simple-minded system of a representation-finite self-injective algebra A can be obtained by irreducible iterative left tilting mutation starting from the set of simple A-modules. In fact, we can already obtain this result using Theorem 4.5 and the following corollary of Dugas' proposition (see [CKL] for more detailed exposition):

Corollary 4.11. *Let $\mathcal{S}$ be a simple-minded system of a representation-finite symmetric algebra A, then there is some simple-minded collection $\mathbf{S}$ with $\eta_A(\mathbf{S}) = \mathcal{S}$. In particular, there is some tilting complex T such that $\mathcal{S}$ is a E_T-simple-image. Moreover, this surjection from tilting complexes to simple-minded systems respects mutations.*

This is our starting point of posing the Problem 1.2, as already mentioned briefly in the introduction.

Proposition 4.12. *Let $\mathcal{S}$ be the set of simple $B^G_{e,m}$-modules, and S_i be a simple $B^G_{e,m}$-module corresponding to an edge i of G. Then an irreducible left mutation $\mu^-_{S_i}(\mathcal{S})$ replaces at most three (indecomposable) modules in $\mathcal{S}$. In particular, at most three (indecomposable) modules in any simple-minded system of any Brauer tree algebras will be replaced after performing an irreducible left mutation.*

Proof. We use the labelling of edges in G appeared in Definition 4.6, also let $\mathcal{S} = \{S_1, \ldots, S_e\}$ with S_x being the simple module corresponding to the edge x in G, and denote P_x as the projective cover of S_x. By the definition of mutation, S_i itself is replaced by $\Omega^{-1}S_i$. For any $S_x \in \mathcal{S}$ with $x \neq i$, there is non-zero (stable) homomorphism from ΩS_x to (if any) self-extensions of S_i, if and only if, the top of $\operatorname{rad} P_x$ contains S_i, which happens if and only if $x = j$ or k (c.f. the labelling in Definition 4.6). One can then explicitly calculate that S_j is replaced by the Loewy length two uniserial $B^G_{e,m}$-module (denoted as $u(S_j, S_i)$) with top S_j and socle S_i; and S_k is replaced by $u(S_k, S_i)$. i.e. $\mu^-_{S_i}(\mathcal{S}) = \{S_n | n \neq i, j, k\} \cup \{\Omega^{-1}S_i, u(S_j, S_i), u(S_k, S_i)\}$. Note that S_j or S_k may not exist depending on the Brauer tree G (more specifically, the valencies of the two ends of edge i). Now suppose $\mathcal{S}$ is an arbitrary simple-minded system of arbitrary Brauer tree algebra, we know from Theorem 2.4 that there is a stable equivalence ϕ making $\mathcal{S}$ a simple-image. The last statement now follows from the fact that $\mu^-_X(\mathcal{S}) = \phi^{-1}(\mu^-_{\phi X}(\phi\mathcal{S}))$. □

Remark 11. Dually, same result holds for irreducible right mutation.

The following observation is crucial for the proof of the main theorem. We will implicitly identify simple-minded systems with configurations to keep the arguments linguistically simple. Also recall from previous section that the type of a configuration indicates the rim where the simple module lies after tree pruning (c.f. Lemma 2.9).

Proposition 4.13. *Let $\mathcal{S}$ is a simple-minded system of $B^{\star}_{e,m}$ which is a $B^{G}_{e,m}$-simple-image for some Brauer tree G, with the valency of exceptional vertex of G being $\ell > 1$. If $X \in \mathcal{S}$ is the image of simple $B^{G}_{e,m}$-module corresponding to an edge attached to the exceptional vertex, then $\mathcal{S}$, $\mu^{+}_{X}(\mathcal{S})$, and $\mu^{-}_{X}(\mathcal{S})$ are all of the same type.*

Proof. Again, stick with the labelling used in the previous proposition (also in Definition 4.6), and suppose X corresponds to edge i attached to exceptional vertex v. We show the proof for left mutation; right mutation result is done analogously. It follows from Proposition 4.12 that the effect of irreducible mutation on the configuration is to replace at most three of the vertices. After tree pruning, the effect on the configuration (corresponding to) $\mathcal{S}$ is $\mathcal{C}_\ell = \{(1, y), \ldots, (\ell, y)\}$ where y is either 1 (bottom-type) or em (top-type). After the mutation at X, all but one modules corresponding to the edges attached to v remain unchanged in $\mu^{-}_{X}(\mathcal{S})$, so after pruning the mutated tree, we are left with a configuration $\mathcal{C}_{\ell-1}$ of $B^{\star}_{\ell-1,m}$ such that the $\ell-2$ vertices in $\mathcal{C}_{\ell-1}$ lie in the same rim as their corresponding vertices in $\omega^{m}_{e}(\mathcal{C}_{\ell-1}) \subset \mathcal{C}_\ell$. When $\ell > 2$, this forces $\mathcal{C}_{\ell-1}$ and $\mathcal{C}_\ell$ to be of the same type, and hence $\mathcal{S}$ and $\mu^{-}_{X}(\mathcal{S})$. If $\ell = 2$, the proposition follows from Example 10 and tree pruning on Brauer line with 2 edges and an exceptional extremal vertex. ☐

5. Proof of Theorem 1.3

This entire section is devoted to prove the main theorem 1.3. Throughout the section, we denote A as the Brauer star algebra $B^{\star}_{e,m}$ with multiplicity $m > 1$; and B as the Brauer star algebra $B^{\star}_{e,1}$. For any algebra C, we denote $\mathcal{S}_C$ as the set of simple C-modules.

As before, we identify the set of simple-minded systems of A with the set of configurations of $\mathbb{Z}A_{me}/\langle\tau^e\rangle$. Now there are maps

$$\begin{tikzcd}
\left\{\begin{array}{c}\text{alternating pointed Brauer trees with}\\ \text{a marked non-exceptional vertex}\end{array}\right\} \arrow[d, leftrightarrow, "f"] & \\
\left\{\begin{array}{c}\text{two-term tilting complexes of } A\\ \text{concentrated in degree } -1,0\end{array}\right\} \arrow[r, "g"] & \left\{\begin{array}{c}\text{configurations of}\\ \mathbb{Z}A_{em}/\langle\tau^e\rangle\end{array}\right\}
\end{tikzcd}$$

where f is a bijection given by the algorithm presented in Section 3, and $g(T) = \overline{F_T}^{-1}(\mathcal{S}_{E_T})$. Note that, using Koenig-Yang bijection and formula (4.1), the map g respect all mutations.

For convenience, we denote the sets above as $\mathrm{AltPt}(e, m)$, 2-tilt(A), and $\mathrm{Con}(A_{em}, 1/m, 1)$ respectively. The elements of $\mathrm{AltPt}(e, m)$ are given by the triple (G, pt, u) where $G = (G, v, m)$ is a Brauer tree with e edges and multiplicity m at vertex v, pt is the choice of alternating pointing (with $pt = even$ for even-first, and $pt = odd$ for odd-first), and u is the marked non-exceptional vertex. Recall that the pointing on Brauer star is unique; however, for technical issues, we redefine instead, that there are two pointings for Brauer star, with the $f(\star, odd, u) = A$, and $f(\star, even, u) = A[1]$ for any u non-exceptional.

Also note that the elements of $\mathrm{AltPt}(e,m)$ are defined up to rotational symmetry of the underlying graph of G around v, i.e. it is possible that $(G,pt,u)=(G,pt,u')$ for two different vertices u,u' in G. In particular, $(\star,pt,u)=(\star,pt,u')$ for all u,u' non-exceptional. We are going to show that the composition gf is a bijection for any $m>1$, and then go on to show that the corresponding map is a surjection when $m=1$.

Lemma 5.1. *For each Ω-orbit in $\mathrm{Con}(A_{em},1/m,1)/\langle\Omega\rangle$, there is a configuration $\mathcal{S}$ in this orbit which is also an image of gf.*

Proof. As discussed in Section 2 Theorem 2.4 there is a bijection:

$$\{\text{isoclasses of Brauer tree algebras with } e \text{ edges and multiplicity } m\}\to \mathrm{Con}(A_{em},1/m,1)/\langle\Omega\rangle$$

Also note that the first set is in bijection with the set of Brauer trees with e edges and multiplicity m, such bijection is given as follows. Suppose G is a Brauer tree and $C=B^{G}_{e,m}$ be its associated Brauer tree algebra, then pick arbitrary stable equivalence $F:\underline{\mathrm{mod}}\text{-}C\to\underline{\mathrm{mod}}\text{-}A$ and configuration (identified with a simple-minded system) $\mathcal{S}=F(\mathcal{S}_C)$. The map then sends C to the Ω-orbit of $\mathcal{S}$. We now refine this map to make it compatible with gf.

As in the proof of Theorem 3.4, let $(G,pt,u)\in\mathrm{AltPt}(e,m)$, we construct the two-term tilting complex T, and get derived equivalence $F_T: D^b(\text{mod-}A)\to D^b(\text{mod-}E_T)$ which induces a stable equivalence $\overline{F_T}:\underline{\mathrm{mod}}\text{-}A\to\underline{\mathrm{mod}}\text{-}E_T$, and so we have $gf(G,pt,u)=\overline{F_T}^{-1}(\mathcal{S}_{E_T})$. Since $\mathrm{AltPt}(e,m)$ contains all possible Brauer trees with e edges and multiplicity m, the assertion follows. □

Lemma 5.2. *If $\mathcal{S}\in\mathrm{Con}(A_{em},1/m,1)$ is in the image of gf, then $\tau^n(\mathcal{S})$ is also an image under gf for all $n\in\mathbb{Z}$.*

Proof. Suppose $\mathcal{S}=gf(G,pt,u)$ for some Brauer tree G with pointing pt and marked vertex u. Now fix G and pt, as we change the marked non-exceptional vertex u, according to algorithm presented in Section 3, we are only changing the labelling on the projective indecomposable modules. More precisely, if we change the marked vertex from u to u' where u' is the (non-exceptional) vertex immediately after u in the Green's walk around the Brauer tree, then we permutes the labelling on the vertices cyclicly (around the Brauer tree), and hence its only effect on the induced two-term tilting complex is a cyclic permutation on the labelling of the projective indecomposable modules. Consequently, the resulting configuration can be obtained by relabelling the x-coordinate of all the vertices in the original one, which corresponds to the effect of applying τ on the original configuration. Since there are e choices of the marked vertex, all the elements in the τ-orbit of $\mathcal{C}$ is then an image under gf. □

To show gf is a bijection, we are left to prove that for any $\mathcal{S}\in\mathrm{Con}(A_{em},1/m,1)$ which is in $\mathrm{Im}(gf)$, then $\Omega^{-1}\mathcal{S}$ is also in $\mathrm{Im}(gf)$. Recall from Section 3 that we divided the two-term star-to-tree tilting complexes into two classes, namely the positive one and negative one. Note that these correspond to even-first pointed and odd-first pointed Brauer tree with marked vertex respectively.

Lemma 5.3. *For any Brauer tree G and a fixed non-exceptional vertex u of G, take $T=f(G,odd,u)$ and $T'=(f(G,even,u))[-1]$, and let $\mathcal{S},\mathcal{S}'$ be the image of simple E_T and $E_{T'}$-module respecitvely. Then $\mathcal{S}$ and $\mathcal{S}'$ are of the same type. In particular, since $\mathcal{S}_A$ is of bottom-type, $\mathcal{S}$ and $\mathcal{S}'$ in the lemma are also of bottom-type.*

Proof. T is a negative tilting complex, so by Theorem 4.8, $T=\mu^-_{P_{i_\ell}}\cdots\mu^-_{P_{i_1}}(A)$. Therefore $\mathcal{S}=g(T)=\overline{F_T}^{-1}(\mathcal{S}_{E_T})$ can be obtained by iterative irreducible left simple-minded system mutations $\mu^-_{X_\ell}\cdots\mu^-_{X_1}(\mathcal{S}_A)$. Apply Proposition 4.13 successively, $\mathcal{S}$ and $\mathcal{S}_A$ is of the same type (bottom-type).

Now let $\mathcal{S}' = \overline{F_{T'}}^{-1}(\mathcal{S}_{E_{T'}})$. Similar to T, since T' is positive, by Theorem 4.8, T can be obtained by ℓ steps irreducible right tilting mutation, inducing ℓ steps irreducible right simple-minded system mutation. Apply Proposition 4.13 successively, $\mathcal{S}'$ and $\mathcal{S}_A$ is of the same type. Hence, $\mathcal{S}'$ and $\mathcal{S}$ are of the same type. □

Lemma 5.4. *If* $\mathcal{S} \in \mathrm{Im}(gf)$ *is a* $B_{e,m}^G$*-simple-image of bottom-type, then there is a* $B_{e,m}^G$*-simple-image of top-type in* $\mathrm{Im}(gf)$*.*

Proof. Suppose $\mathcal{S} = gf(G, odd, u)$ for some u, and let $U = f(G, even, u) \in$ 2-tilt$A$ and $T' = U[-1]$ as in the statement of the previous lemma. Obviously $\mathrm{End}_{\mathcal{T}}(T') = \mathrm{End}_{\mathcal{T}}(U)$, and so $\mathcal{S}_{E_{T'}} \cong \mathcal{S}_{E_U}$. Now we have

$$\begin{aligned}
\mathcal{S}' &= \overline{F_{T'}}^{-1}(\mathcal{S}_{E_{T'}}) \\
&= \overline{F_{U[-1]}}^{-1}(\mathcal{S}_{E_{U[-1]}}) \\
&= \overline{\mathrm{Hom}(U[-1], -)}^{-1}(\mathcal{S}_{E_U}) \\
&= \overline{\mathrm{Hom}(U, -) \circ [1]}^{-1}(\mathcal{S}_{E_U}) \\
&= (\overline{F_U} \circ \Omega^{-1})^{-1}(\mathcal{S}_{E_U}) \\
&= \Omega(\overline{F_U}^{-1}(\mathcal{S}_{E_U})) \\
&= \Omega(gf(G, even, u))
\end{aligned}$$

The previous lemma says that $\mathcal{S}'$ is of bottom-type, and so $gf(G, even, u) = \Omega^{-1}\mathcal{S}'$ is of top-type. □

Finally, note that if $\mathcal{S}$ is of bottom-type, then $\Omega^{-1}\mathcal{S}$ is of top-type. Applying Lemma 5.2 to the above lemma, we have that every top-type and bottom-type $B_{e,m}^G$-simple-image is in $\mathrm{Im}(gf)$. This completes the proof for the bijection part of the main theorem.

Lemma 5.5. *We have the bijection* 2-tilt$(A) \leftrightarrow$ 2-tilt(B)*.*

Proof. Note that in given a Brauer tree with multiplicity 1, we can just pick any vertex to be an "exceptional vertex". Now the statement follows from the fact that AltPt(e, m)=AltPt$(e, 1)$ and Schaps-Zakay-Illouz correspondence (Theorem 3.3). □

Suppose G, G' are a Brauer trees of multiplicity 1 with the same underlying tree and "exceptional vertex" v, v' respectively. Let u be a vertex in G (and G') which is not v and v', and $T = f(G, pt, u), T' = f(G', pt, u)$. Now by Lemma 5.5 and Schaps-Zakay-Illouz correspondence $E_T \cong B_{e,1}^G \cong B_{e,1}^{G'} \cong E_{T'}$, and so $g(T)$ and $g(T')$ are in the same Ω-orbit. Now surjectivity of the map g follows by applying all the previous lemmas analogously. We have now completed the proof of the main theorem.

Acknowledgement The author would like to thank Alexandra Zvonareva for directing the attention to two-term tilting complexes via work of [AH] and sharing her calculations, which lead to the creation of this work. The author is supported by EPSRC Doctoral Training Scheme.

References

[Ada] T. Adachi. τ-tilting modules of Nakayama algebras. 2013. In Preparation.

[AH] H. Abe and M. Hoshino. On derived equivalences for self-injective algebras. *Comm. Algebra*, 34(12):4441–4452, 2006.

[AI] T. Aihara and O. Iyama. Silting mutation in triangulated categories. *J. Lond. Math. Soc. (2)*, 85(3):633–668, 2012.

[Aih1] T. Aihara. Mutating Brauer trees. 2010. `arXiv:1009.3210`.

[Aih2] T. Aihara. Tilting-connected symmetric algebras. *Algebr Represent Theor.* DOI: 10.1007/s10468-012-9337-3. 2012.

[AIR] T. Adachi, O. Iyama, and I. Reiten. τ-tilting theory. 2012. `arXiv:1210.1036`.

[AN] S. Al-Nofayee. Equivalences of derived categories for selfinjective algebras. *J. Algebra*, 313(2):897–904, 2007.

[AZ] M. Antipov and A. Zvonareva. Two-term partial tilting complexes over Brauer tree algebras. 2013. `arXiv:1302.3342`.

[CKL] A. Chan, S. Koenig, and Y.M. Liu. Simple-minded systems, configurations and mutations of representation-finite selfinjective algebras. 2013. In Preparation.

[Dug1] A. Dugas. Tilting mutation of weakly symmetric algebras and stable equivalence. 2011. `arXiv:1110.1679`.

[Dug2] A. Dugas. Torsion pairs and simple-minded systems in triangulated categories. 2012. `arXiv:1207.7338`.

[GR] P. Gabriel and Ch. Riedtmann. Group representations without groups. *Comment. Math. Helvetici*, 54:240–287, 1979.

[Gre] J. A. Green. Walking around the Brauer Tree. *J. Austral. Math. Soc.*, 17:197–213, 1974.

[KL] S. Koenig and Y. Liu. Simple-minded systems in stable module categories. *Quart. J. Math.*, 2011.

[KY] S. Koenig and D. Yang. Silting objects, simple-minded collections, t-structure and co-t-structure of finite dimensional algebras. 2012. `arXiv:1203.5657`.

[Miz] Y. Mizuno. ν-stable τ-tilting modules. 2012. `arXiv:1210.8322`.

[Ric1] J. Rickard. Derived categories and stable equivalence. *J. Pure Appl. Algebra*, 61(3):303–317, 1989.

[Ric2] J. Rickard. Morita theory for derived categories. *J. London Math. Soc.*, 39:436–456, 1989.

[Rie1] Ch. Riedtmann. Algebren darstellungskocher, uberlargerungen and zurick. *Comment. Math. Helv.*, 55(2):199 – 224, 1980.

[Rie2] Ch. Riedtmann. Representation-finite selfinjective algebras of class A_n. *Lecture Notes in Math. Springer*, 832:449 – 520, 1980.

[Rie3] Ch. Riedtmann. Representation-finite selfinjective algebras of class D_n. *Compositio Mathematica*, 49(2):231–282, 1983.

[RS] J. Rickard and M. Schaps. Folded tilting complexes for Brauer tree algebras. *Adv. Math.*, 171(2):169–182, 2002.

[SZI] M. Schaps and E. Zakay-Illouz. Pointed Brauer trees. *J. Algebra*, 246(2):647–672, 2001.

Aaron Chan
Institute of Mathematics
University of Aberdeen
Aberdeen, AB24 3UE
UK

E-mail address: `aaron.kychan@gmail.com`